Chem 11	D Fall 2016	Exam III	Whela	
SID	Last Key	First Ans	swer	
Question 1	a. Write a net ionic equation to sho	ow that hydrosulfuric acid,	behaves as an acid in	
6 Points	water. $H_2S(aq) + H_2O(I)$	H ₃ 0 ⁺ +	HS-	
	(= or	⇔)		
	 b. Write a net ionic equation to sho water. 	ow how sodium hydroxide b	ehaves as a base in	
		Nat	OH-	
	NaOH(aq) (= or	⇔)		
Question 2 8 Points	a. HNO3	1. Strong Acid		
	ь. нсоон <u>2</u>	2. Weak Acid		
	c. C₅H₅N <u>4</u>	3. Strong Base		
	d. NH₄⁺2	4 . Weak Base		
Question 3 4 Points	An aqueous solution has a hydroxide ion	concentration of 1.0 X 10) ⁻² M.	
	a) What is the hydronium ion concentration in this solution?			
	b) Is this solution acidic , basic or n	eutral?	Basic	
Question 4 6 Points	An aqueous solution has a pOH of 6			
	a) What is the pH of this solution?		8	
	b) What is the hydronium ion conce	ntration in this solution?	<u>IXI0</u> M	
	c) What is the hydroxide ion conce	ntration in this solution?	<u> X 10⁻⁶</u> M	
Question 5 6 Points	Arrange the following solutions in order 1 = least acidic ; 3 = most acidic	of increasing acidity:		
	a) Solution with a pH = 11		1	
	b) Solution with a hydroxide ion cor	centration = 1×10^{-11} M	3	
	c) Solution with a hydronium ion con	3 2		
	•			
Question 6 6 Points	The hydronium concentration in an aque			
	a. The hydroxide ion concentration	<u>.</u>		
	b. The pH of this solution is:	<u>1.45</u>		
	c. The pOH is:	12.55		

Question 7	a) For following net ionic equation:			
	HClO(aq) + H₂O(l) ⇔ ClO ⁻ + H₃O ⁺ - Circle the appropriate answer - B-L = Bronsted Lowry			
	H ₂ O B-L Acid B-L Base			
	CIO ⁻ B-L Acid B-L Base			
	b) The formula for the conjugate <u>Base</u> of H_3O^+ is: <u>H_2O^-</u>			
	c) The formula for the conjugate Ocid of CIO is:			
Question 8	A buffer solution that is 0.436M in HCN and 0.436M in KCN has a pH of 9.40 .			
	Addition of which of the following would increase the capacity of the buffer for add H_3O^2 ?			
	🖒 KCN 🗆 HCN			
	🕞 both HCN and KCN 🛛 pure water			
	\Box none of these choices			
Question 9	Which of the following aqueous solutions are buffer solutions ?			
	 ○ 0.14M HF + 0.17M KF □ 0.34M Ba(ClO₄)₂ + 0.25M BaI₂ 			
	□ 0.19M Ca(OH) ₂ + 0.21M CaCl ₂ ○ 0.34M NH ₄ NO ₃ + 0.34M NH ₃			
	□ 0.25M HCl + 0.17M KCl			
Question 10 6 Points	A buffer solution is made that is 0.472M in H_2CO_3 and 0.472M in NaHCO ₃ .			
	a) Ka for H_2CO_3 is 4.2×10 ⁻⁷ , what is the pH of the buffer solution? 6.4			
	b) Write the net ionic equation for the reaction that occurs when 0.129 mol NaOH is added to 1.00 L of the buffer solution.			
	$H_{2}CO_{3}(a_{1}) + OH^{-} = H_{2}O(l) + H_{2}O_{3}$			
Question 11	A buffer solution is 0.440 M in HCN and 0.324 M in NaCN. If Ka for HCN is 4.0x10 ⁻¹⁰ ,			
6 Points	what is the pH of this buffer solution?			
	Must show work $pH = -\log_{10}(4.0 \times 10^{-10}) + \log_{10}\frac{[CN]}{[HCN]}$			
	$= 9.397 + \log_{10}\left(\frac{0.324}{0.440}\right)$			
	$= 9.397 + \log_{10}(0.736)$			
	= 9.397 - 0.133			
	pH = 9.16			

Question 12 6 Points	A small amount of strong acid is added to a buffer made from HCN and NaCN . What changes if any will occur to the following.			
	Choose from the following choices: Increase significantly Increase Increase slightly Decrease significantly Decrease Decrease slightly			
	a) pOH <u>Americase slightly</u>			
	b) [HCN]			
Question 13 6 Points	The isotope ⁶⁰ 27 Co is but one of many isotopes whose Neutron/Proton ratio is too large .			
	a) The only form of radioactive decay available to ⁶⁰ 27Co is: <u>Belo effission</u>			
	b) The balanced nuclear equation for this decay: ⁶⁰ 27Co = <u>-1C</u> +28N1			
Question 14 6 Points	Write a balanced nuclear equation for the following:			
o roints	a) $^{214}_{82}$ Pb undergoing beta decay: $\frac{214}{280} = \frac{-12}{280} = \frac{-12}{280}$			
	b) ${}^{28}{}_{15}P$ undergoing positron emission: $4 - 10^{28}$ = $\frac{+10^{28} + 14^{28}}{4 - 10^{28}}$			
	c) ⁴¹ 20 Ca undergoing electron capture: <u>20(a + 1)</u> = <u>19 K</u>			
Question 15 6 Points	How many moles of water will be formed upon the complete reaction of 27.3 grams of sulfuric acid with excess zinc(II) hydroxide ? sulfuric acid (aq) + zinc(II) hydroxide (s) = zinc(II) sulfate (aq) + water (l) For full credit you must show work and include a balanced chemical equation.			
	$H_{2}SO_{4}(aq) + Z_{1}(oH)_{2}(aq) = Z_{1}SO_{4}(aq) + 2H_{2}O(1)$			
	27.3g			
	$H_{2}50_{4}: 2(1.01) + 32.07 + 4(16.00) = 98.09 g.mol^{-1}$			
	$27.39 H_2 504 t mol = 0.278 mol H_2 504 98.099 = 0.278 mol H_2 504$			
	$0.278 \mod H_2 SO_4 2 \ H_2 D = 0.557 \mod H_2 O$ $1 \ H_2 SO_4$			
	0.577 mol			

An aqueous solution of **barium hydroxide** is standardized by titration with a **0.140 M** Question 16 8 Points solution of hydrochloric acid. If 26.8 mL of base are required to neutralize 19.4 mL of the acid, what is the molarity of the barium hydroxide solution? For full credit you must show work and include a balanced chemical equation. $Be(OH)_2 + 2HOP = BeOl_2 + 2H_2O$ 26.8 mL 19.4 mL 0.140 M #mol Hep = 0.140 (0.0194) = 2.72 × 10-3 mol Hep $2.72 \times 10^{-3} \text{ mol Her} | 1 \text{ Ba(oH)}_{2} = 1.36 \times 10^{-3} \text{ mol Ba(oH)}_{2}$ $M = \frac{1.36 \times 10^{-3}}{0.0268} = 0.0507$ 0.0507 M Question 17 What volume of a 0.142 M solution of aluminum bromide contains the same number of 6 Points moles of **aluminum bromide** as there are in **43.2 mL** of a **0.124 M** solution of **aluminum** bromide? Must show work $\# \text{ mol alBr}_{3} = 0.124(0.0432) = 5.36 \times 10^{-3}$ # mol OPBr3 = M X V(L) 5.36 × 10-3 = 0.142 × 1(L) $V(L) = \frac{5.36710^{-3}}{0.142} = 0.0377$ 0.03 L

Exam III Score