

Announcements – Lecture XXIII – Wednesday, June 24th

1. Exam III:

Friday, June 26th, In Class

3 or 4 questions will be taken from Lab Owls:-


3.4 , 4.2 , 4.5 , 5.5 , 5.6



10.5 Kinetic Molecular Theory

The Postulates

- 1) The volume occupied by the gas molecules is negligible in comparison to the volume of the container they are in.
- 2) Collisions between gas molecules are totally elastic ... no loss of energy ... no intermolecular force of attraction
- 3) \bar{KE} is proportional to temperature ... at a given temperature all gases have the same average kinetic energy (\bar{KE}), regardless of their mass.




PRESSURE = Force per unit area

- a) Energy of the collisions with the walls of the container.
- b) The frequency of these collisions.


10.5 Kinetic Molecular Theory

A: Kinetic Molecular Theory and the Gas Laws



10.5 Kinetic Molecular Theory

A: Kinetic Molecular Theory and the Gas Laws – Charles's Law



n and V are held constant

Increase T , what happens to P ?

$P \uparrow$... Why?

- a) Collisions MORE energetic ... $\overline{KE} \uparrow$
- b) Collision frequency increases.


$$P \propto T$$

$$\frac{P}{T} = \text{constant}$$

$$\frac{P_1}{T_1} = \frac{P_2}{T_2} \dots \text{Charles's law}$$

10.5 Kinetic Molecular Theory

A: Kinetic Molecular Theory and the Gas Laws – Avogadro's Law



V and T are held constant

Increase n, what happens to P?

P↑ ... why?

With more particles the frequency of the collisions increases.


$$P \propto n$$

$$\frac{P}{n} = \text{constant}$$

$$\frac{P_1}{n_2} = \frac{P_2}{n_1} \dots \text{Avogadro's Law}$$

10.5 Kinetic Molecular Theory

A: Kinetic Molecular Theory and the Gas Laws – Boyle's Law



n and T are held constant.

$$P \propto \frac{1}{V}$$

Decrease V , what happens to P ?

PT ... why?

In a smaller volume the frequency of the collisions increases.

$$PV = \text{constant}$$

$$P_1V_1 = P_2V_2 \dots \text{Boyle's Law}$$

10.5 Kinetic Molecular Theory

A: Kinetic Molecular Theory and the Gas Laws – Ideal Gas Law

$$P \propto \frac{nT}{V}$$

$$\frac{PV}{nT} = \text{Constant} = R = \text{Ideal Gas Constant}$$

$8.314 \text{ J.mol}^{-1}\text{.K}^{-1}$

$0.0821 \text{ L.atm.K}^{-1}\text{.mol}^{-1}$

$$\frac{PV}{nT} = R$$

$$PV = nRT \rightarrow \text{Ideal Gas Law}$$

10.5 Kinetic Molecular Theory

Memorizing all the Gas Laws ... ☺

GENERAL GAS LAW : n held constant

$$PV = nRT$$

$$\frac{PV}{T} = nR \dots \text{Constant}$$

$$\frac{P_1V_1}{T_1} = \frac{P_2V_2}{T_2}$$

YOUR TURN ... pick two from P, V, n, T ... these to be held constant ... derive the associated Gas Law.

10.5 Kinetic Molecular Theory

D: Nonideal Gases

- a) Gas has Negligible volume
 - b) No intermolecular forces
- Want $PV = nRT$ to work
- keep the pressure low.
→ keep the temperature high.

van der Waals

$$(P + a \left[\frac{n}{V} \right]^2) (V - b n) = nRT$$

↑
Intermolecular forces correction

Gas molecule volume correction.