



14.5 Activation Energy and Temperature

Reaction Coordinate Diagrams



E_a = Activation Energy.

$$\Delta E = E_{\text{PRODUCTS}} - E_{\text{REACTANTS}}$$

< 0
Exothermic

$$\Delta E = E_{\text{PRODUCTS}} - E_{\text{REACTANTS}}$$

> 0
Endothermic

14.5 Activation Energy and Temperature

The Arrhenius Equation

$$k = A e^{-\frac{E_0}{RT}}$$

k : Rate constant.

A : Frequency factor.

E_0 : Activation energy.

R : Ideal Gas Constant.

T : Temperature in K.

A : Measure of the number of collisions that take place with the correct orientation.

$e^{-\frac{E_0}{RT}}$: Fraction of the collisions that occur with sufficient energy to overcome E_0 .

a) $E_0 \uparrow$, then $e^{-\frac{E_0}{RT}} \downarrow$, $k \downarrow$

b) $E_0 \downarrow$, then $e^{-\frac{E_0}{RT}} \uparrow$, $k \uparrow$

$$k_1 = A e^{-\frac{E_0}{RT_1}} \quad : \quad k_2 = A e^{-\frac{E_0}{RT_2}}$$

$$\ln k_1 = \ln \left(A e^{-\frac{E_0}{RT_1}} \right) \quad : \quad \ln k_2 = \ln \left(A e^{-\frac{E_0}{RT_2}} \right)$$

$$\ln k_2 - \ln k_1 = \ln A + \ln e^{-\frac{E_0}{RT_2}} - \ln A - \ln e^{-\frac{E_0}{RT_1}}$$


$$\ln k_2 - \ln k_1 = -\frac{E_0}{RT_2} - \left(-\frac{E_0}{RT_1} \right)$$

$$\ln \frac{k_2}{k_1} = -\frac{E_0}{R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right)$$

14.5 Activation Energy and Temperature

The Arrhenius Equation

The **activation energy** for the gas phase decomposition of acetic anhydride is 144 kJ/mol.



The **rate constant** for this reaction is $6.02 \times 10^{-4} \text{ s}^{-1}$ at 495 K. What is the **rate constant** at 531 K?

$$\ln \frac{k_2}{k_1} = -\frac{E_0}{R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right)$$

$$E_0 = 144,000 \text{ J} \quad (\text{Remember, } R \text{ is in J})$$

$$T_1 = 495 \text{ K} \quad T_2 = 531 \text{ K}$$

$$k_1 = 6.02 \times 10^{-4} \quad k_2 = ?$$

$$R = 8.314 \text{ J mol}^{-1} \text{ K}^{-1}$$

$$\ln \frac{k_2}{6.02 \times 10^{-4}} = -\frac{144,000}{8.314} \left(\frac{1}{531} - \frac{1}{495} \right)$$

$$\ln k_2 - \ln(6.02 \times 10^{-4}) = -17320.2 (-1.3696 \times 10^{-4})$$

$$\ln k_2 + 7.4152 = 2.3722$$

$$\ln k_2 = 2.3722 - 7.4152$$

$$\ln k_2 = -5.043$$

$$k_2 = 6.45 \times 10^{-3} \text{ s}^{-1}$$

14.5 Activation Energy and Temperature

Graphical Determination of E_a

$$k = Ae^{-\frac{E_a}{RT}}$$
$$\ln k = \ln A + \ln e^{-\frac{E_a}{RT}}$$

$$\ln k = -\frac{E_a}{RT} + \ln A$$

$$\ln k = -\frac{E_a}{R} \left(\frac{1}{T} \right) + \ln A$$


$$y = mx + c$$

$$\ln k \text{ vs } 1/T \text{ plot, slope} = -\frac{E_a}{R}$$

The Arrhenius Equation

Question 1 of 3

The rate of the reaction




is measured at different temperatures, with the following rate constants, k , determined:

Temperature, K	k , s^{-1}
298	3.46×10^{-5}
328	1.5×10^{-3}
358	3.34×10^{-2}
378	0.21

What is the activation energy, E_a , for this reaction in units of kilojoules?

Submit

- k
- $\ln k$
- $1/k$



Least Squares Analysis

- T
- $\ln T$
- $1/T$

Enter a response, then press **SUBMIT**.

kJ

Plot

Clear

14.5 Activation Energy and Temperature

Graphical Determination of E_a

$$\text{slope} = -1.23 \times 10^4$$

$$-\frac{E_a}{R} = -1.23 \times 10^4$$

$$E_a = 1.23 \times 10^4 (8.314)$$


$$E_a = 1.02 \times 10^5 \text{ J. mol}^{-1}$$

OR
 102 kJ. mol^{-1}

The Arrhenius Equation

Question 1 of 3

The rate of the reaction




is measured at different temperatures, with the following rate constants, k , determined:

Temperature, K	k , s^{-1}
298	3.46×10^{-5}
328	1.5×10^{-3}
358	3.34×10^{-2}
378	0.21

What is the activation energy, E_a , for this reaction in units of kilojoules?

Submit

- k
 $\ln k$
 $1/k$



- T $\ln T$ $1/T$

Least Squares Analysis

Enter a response, then press **SUBMIT**.

Plot

102 kJ

Clear

14.6 Reaction Mechanisms and Catalysis

The Components of a Reaction Mechanism

Chemistry Interactive: Mechanism of The Reaction Between NO_2 and CO

Mechanism 1

▶ **Step 1.** (slow, rate-determining step)
 $\text{NO}_2(\text{g}) + \text{CO}(\text{g}) \rightarrow \text{NO}(\text{g}) + \text{CO}_2(\text{g})$



Why is the Initial Rate $\propto k[\text{NO}_2][\text{CO}]$?

BECAUSE experimentally it was found to be



How to explain this ? ... Mechanisms.

Mechanism 2, Step 1

▶ **Step 1.** (slow, rate-determining step)
 $2 \text{NO}_2(\text{g}) \rightarrow \text{NO}_3(\text{g}) + \text{NO}(\text{g})$



Mechanism 2, Step 2

▶ **Step 2.** (fast)
 $\text{NO}_3(\text{g}) + \text{CO}(\text{g}) \rightarrow \text{NO}_2(\text{g}) + \text{CO}_2(\text{g})$

